

Analytical chemistry year 12

Name _____

- 1) Consider the ¹H NMR spectrum on the right.
 - a) Draw the grouping of atoms that would give rise to the triplet and quartet splitting patterns.
 - b) What group of atoms will most likely give rise to the singlet?




- c) Propanol(C_3H_8O) has two isomers. Draw the two possible structures.
- d) Which isomer is represented by the ¹H NMR on the right?



4 marks

- 2) An organic compound has the following percentage composition by mass. 54.55%C, 9.10%H, 36.35%O
 - a) Find its molecular formula if 0.500 mol of the substance weighs 44.0 grams 2 marks
 - b) The IR spectrum of the compound is shown on the right?Draw two possible structures for this compound




i)

ii)


iii) Stephen suggested that the compound might be butanoic acid. Is he right?


Explain with reference to the IR spectrum.

 c) The compound was found not to react with a base and its ¹H NMR is shown on the right.
Draw the possible structure of the molecule.



d) Jonathon was given the ¹H NMR shown below and came up with the same structure as in c) above. Is he correct or incorrect. Explain





3) Two paper chromatograms are shown on the right of a sample of food dye. Do they have a common component? Explain why

2 marks

4) A student is to accurately determine the concentration of a solution of sodium hydrogencarbonate in a titration against a standard solution of hydrochloric acid, HCl.

The first step in this experiment is to accurately dilute 50.0 mL of a 1.00 M HCl stock solution to a 0.100 M solution using a 500 mL volumetric flask. However, instead of using distilled water in the dilution, the student mistakenly adds 100.0 mL of 0.0222 M NaOH, solution before feeling to the mark with distilled..

a) Write an equation for the reaction that occurs in the 500 mL volumetric flask.

2 marks

b) Calculate the <u>final</u> concentration of the hydrochloric acid in the 500mL volumetric flask. Give your answer to correct significant figures.

2 marks

The student then uses this contaminated hydrochloric acid solution to determine the accurate concentration of an unknown sodium hydrogen carbonate (NaHCO₃) solution. During the titration the NaHCO₃ solution is placed in the burette while a 20.0 mL aliquot of the HCl solution is placed in the conical flask

c) Will the calculated concentration of sodium hydrogen carbonate solution be greater or smaller than the true value? Justify your answer?

1 mark

- 5) A mixture of organic compounds is run through a chromatography column. The stationary phase is coated with a non-polar substance while the mobile phase is a polar solvent.
 - a) Place the molecules on the right in order of increasing retention time.

i)

iv)

High

b) Explain your reasons why you placed the

molecules in the order given in a).

2 marks