## Thermochemical equations Lesson 1

 $2C_8H_{18}(g)+25O_2(g) => 16CO_2(g)+18H_2O(g) \ \Delta H = -10900 \ kJ \ molWhat does a thermochemical equation reveal about a reaction?$ 

i)	What is enthalpy?
ii)	What is enthalpy change (ΔΗ)?
iii)	Stoichiometric ratio of a balanced chemical equation
iv)	States
v)	Energy output or input (depending on sign) negative sign indicates
	positive sign indicates
vi)	Amount of energy is dependent on the mol of the equation  For example
1)	Consider the thermochemical equation for the combustion of octane shown below. $2C_8H_{18}(g) + 25O_2(g) \Rightarrow 16CO_2(g) + 18H_2O(g)$ $\Delta H = -10900$ kJ moli. It shows that for 2 mol of octane that reacts 10900 kJ of energy is ii. Calculate the $\Delta H$ for the following equations
	a) $C_8H_{18}(g) + 12\frac{1}{2}O_2(g) => 8CO_2(g) + 9H_2O(g) \Delta H = ? kJ mol^{-1}$

o) $8CO_2(g) + 9H_2O(g) => C_8H_{18}(g) + 12½O_2(g) \Delta H = ? kJ mol$			
) Cons	sider the equation below		
C <sub>6</sub> H <sub>12</sub> O	$_{6}(aq) + 6O_{2}(g) => 6CO_{2}(g) + 6H_{2}O(I) \Delta H = -2803 \text{ kJ mol}^{-}$		
Vhat i	s the ΔH for the following equations?		
	i. $2C_6H_{12}O_6(aq) + 12O_2(g) \Rightarrow 12CO_2(g) + 12H_2O(I) \Delta H =$		
	ii. $3CO_2(g) + 3H_2O(I) => \frac{1}{2} C_6H_{12}O_6(aq) + 3O_2(g) \Delta H =$		
<sub>6</sub> H <sub>12</sub> O	<sub>6</sub> (aq) + 6O <sub>2</sub> (g) => 6CO <sub>2</sub> (g) + 6H <sub>2</sub> O(I) ΔH = -2803 kJ mol <sup>-</sup>		
-	et amount of energy, in kJ, is released if 8.80 grams of carbon dioxide is sed when glucose burns in excess oxygen?		

g) Wha	t amount of energy, in kJ, is released if 3.60 grams of glucose reacts
comple	etely?
	<del></del>
	sider the equation below.
	sider the equation below. ) + $6H_2O(I) \Rightarrow C_6H_{12}O_6(aq) + 6O_2(g)$
	) + $6H_2O(I) => C_6H_{12}O_6(aq) + 6O_2(g)$
	) + $6H_2O(I) => C_6H_{12}O_6(aq) + 6O_2(g)$
	) + $6H_2O(I) => C_6H_{12}O_6(aq) + 6O_2(g)$
	) + $6H_2O(I)$ => $C_6H_{12}O_6(aq)$ + $6O_2(g)$ i. What is the $\Delta H$ of the thermochemical equation above ?
	) + $6H_2O(I)$ => $C_6H_{12}O_6(aq)$ + $6O_2(g)$ i. What is the $\Delta H$ of the thermochemical equation above ?  ii. Is energy absorbed or released?
	i. What is the $\Delta H$ of the thermochemical equation above ? ii. Is energy absorbed or released? iii. What amount of energy, is involved if 36.0 grams of glucose (mo
	) + $6H_2O(I)$ => $C_6H_{12}O_6(aq)$ + $6O_2(g)$ i. What is the $\Delta H$ of the thermochemical equation above ?  ii. Is energy absorbed or released?
	i. What is the $\Delta H$ of the thermochemical equation above ? ii. Is energy absorbed or released? iii. What amount of energy, is involved if 36.0 grams of glucose (mo
	i. What is the $\Delta H$ of the thermochemical equation above ? ii. Is energy absorbed or released? iii. What amount of energy, is involved if 36.0 grams of glucose (mo
	i. What is the $\Delta H$ of the thermochemical equation above ? ii. Is energy absorbed or released? iii. What amount of energy, is involved if 36.0 grams of glucose (mo