




Quiz 6 – revision of redox and galvanic cells.

1. Consider the galvanic cell shown on the right.
 - a. What is the theoretical EMF produced by the cell if it is run at standard conditions.
 - b. Write the half equations to each half-cell in the space provided.
 - c. On the diagram label the:
 - i. Direction of electron flow.
 - ii. Direction of positive ions flow from the salt bridge
 - iii. Anode and give its polarity
 - iv. Cathode and give its polarity.
 - d. What will happen to the mass of each electrode as the galvanic cell is allowed to operate. Explain you answer.
 - e. What material should electrode “A” be made from?






2. Below are two redox equations. For each equation identify the :
 - atom being reduced (justify your answer using oxidation numbers)
 - atom being oxidised (justify your answer using oxidation numbers)
 - oxidant
 - reductant



3. Give the oxidation number of underlined atoms in each of the following substances.
 - a. MnO₄
 - b. H₂O₂
 - c. Mg(OH)₂
 - d. CrO₄
 - e. C₃H₈O

4. Consider the following unbalanced redox reactions. Write balanced half equations for each, states not included, and identify each one as either an oxidation or reduction.



5. Using the E° series shown on the right predict if a spontaneous reaction will occur when:

i. Tin(Sn) metal is placed in a 0.1 M AgNO_3 solution.

ii. Pure manganese metal is placed in a solution of 0.2M $\text{Al}(\text{NO}_3)_3$

iii. Nickel metal is placed in a solution of $\text{Fe}(\text{NO}_3)_3$.

iv. Lithium metal is placed in a 1.0 M solution of $\text{Zn}(\text{NO}_3)_2$.

v. Iron metal is placed in a 0.10 M HCl solution.

vi. Magnesium metal is placed in a 1.0M HCl solution.

6. For each spontaneous reaction that occurs in question 5, above, give the

i. oxidant and reductant taking part in the reaction

ii. Write the balanced half equations, with states

iii. Write the balanced overall equation, with states.

Reaction	Standard electrode (E°) in volts at :
$\text{F}_2(\text{g}) + 2\text{e}^- \rightleftharpoons 2\text{F}^-(\text{aq})$	+2.87
$\text{H}_2\text{O}_2(\text{aq}) + 2\text{H}^+(\text{aq}) + 2\text{e}^- \rightleftharpoons 2\text{H}_2\text{O}(\text{l})$	+1.77
$\text{Au}^+(\text{aq}) + \text{e}^- \rightleftharpoons \text{Au}(\text{s})$	+1.68
$\text{Cl}_2(\text{g}) + 2\text{e}^- \rightleftharpoons 2\text{Cl}^-(\text{aq})$	+1.36
$\text{O}_2(\text{g}) + 4\text{H}^+(\text{aq}) + 4\text{e}^- \rightleftharpoons 2\text{H}_2\text{O}(\text{l})$	+1.23
$\text{Br}_2(\text{l}) + 2\text{e}^- \rightleftharpoons 2\text{Br}^-(\text{aq})$	+1.09
$\text{Ag}^+(\text{aq}) + \text{e}^- \rightleftharpoons \text{Ag}(\text{s})$	+0.80
$\text{Fe}^{3+}(\text{aq}) + \text{e}^- \rightleftharpoons \text{Fe}^{2+}(\text{aq})$	+0.77
$\text{O}_2(\text{g}) + 2\text{H}^+(\text{aq}) + 2\text{e}^- \rightleftharpoons \text{H}_2\text{O}_2(\text{aq})$	+0.68
$\text{I}_2(\text{s}) + 2\text{e}^- \rightleftharpoons 2\text{I}^-(\text{aq})$	+0.54
$\text{O}_2(\text{g}) + 2\text{H}_2\text{O}(\text{l}) + 4\text{e}^- \rightleftharpoons 4\text{OH}^-(\text{aq})$	+0.40
$\text{Cu}^{2+}(\text{aq}) + 2\text{e}^- \rightleftharpoons \text{Cu}(\text{s})$	+0.34
$\text{Sn}^{4+}(\text{aq}) + 2\text{e}^- \rightleftharpoons \text{Sn}^{2+}(\text{aq})$	+0.15
$\text{S}(\text{s}) + 2\text{H}^+(\text{aq}) + 2\text{e}^- \rightleftharpoons \text{H}_2\text{S}(\text{g})$	+0.14
$2\text{H}^+(\text{aq}) + 2\text{e}^- \rightleftharpoons \text{H}_2(\text{g})$	0.00
$\text{Pb}^{2+}(\text{aq}) + 2\text{e}^- \rightleftharpoons \text{Pb}(\text{s})$	-0.13
$\text{Sn}^{2+}(\text{aq}) + 2\text{e}^- \rightleftharpoons \text{Sn}(\text{s})$	-0.14
$\text{Ni}^{2+}(\text{aq}) + 2\text{e}^- \rightleftharpoons \text{Ni}(\text{s})$	-0.23
$\text{Co}^{2+}(\text{aq}) + 2\text{e}^- \rightleftharpoons \text{Co}(\text{s})$	-0.28
$\text{Fe}^{2+}(\text{aq}) + 2\text{e}^- \rightleftharpoons \text{Fe}(\text{s})$	-0.44
$\text{Zn}^{2+}(\text{aq}) + 2\text{e}^- \rightleftharpoons \text{Zn}(\text{s})$	-0.76
$2\text{H}_2\text{O}(\text{l}) + 2\text{e}^- \rightleftharpoons \text{H}_2(\text{g}) + 2\text{OH}^-(\text{aq})$	-0.83
$\text{Mn}^{2+}(\text{aq}) + 2\text{e}^- \rightleftharpoons \text{Mn}(\text{s})$	-1.03
$\text{Al}^{3+}(\text{aq}) + 3\text{e}^- \rightleftharpoons \text{Al}(\text{s})$	-1.67
$\text{Mg}^{2+}(\text{aq}) + 2\text{e}^- \rightleftharpoons \text{Mg}(\text{s})$	-2.34
$\text{Na}^+(\text{aq}) + \text{e}^- \rightleftharpoons \text{Na}(\text{s})$	-2.71
$\text{Ca}^{2+}(\text{aq}) + 2\text{e}^- \rightleftharpoons \text{Ca}(\text{s})$	-2.87
$\text{K}^+(\text{aq}) + \text{e}^- \rightleftharpoons \text{K}(\text{s})$	-2.93
$\text{Li}^+(\text{aq}) + \text{e}^- \rightleftharpoons \text{Li}(\text{s})$	-3.02